

論文全著者名. 論題. 雑誌名. 発行年; 巻 :始頁-終頁	著者名
上浪 健、木村紀久、森 雅秀: 腫瘍随伴症候群としてのTrousseau症候群. 呼吸と循環 2014; 62(12): 1195-1200.	上浪 健、 森 雅秀
上浪 健、森 雅秀、揚塩文崇、香川浩之、松井秀記、里見明俊、新中 学、木村裕美、矢野幸洋、米田 勉、山口俊彦、横田総一郎. Docetaxel投与で過敏反応を起こしたがnab-paclitaxel投与は可能であった肺扁平上皮癌の1例. 癌と化学療法 2015; 42(3):375-377.	上浪 健 森 雅秀 里見明俊 新中 学 木村裕美 矢野幸洋 米田 勉 山口俊彦 横田総一郎
Oji Y, Tatsumi N, Fukuda M, Nakatsuka S, Aoyagi S, Hirata E, Nanchi I, Fujiki F, Nakajima H, Yamamoto Y, Shibata S, Nakamura M, Hasegawa K, Takagi S, Fukuda I, Hoshikawa T, Murakami Y, <u>Mori M</u> , Inoue M, Naka T, Tomonaga T, Shimizu Y, Nakagawa M, Hasegawa J, Nezu R, Inohara H, Izumoto S, Nonomura N, Yoshimine T, Okumura M, Morii E, Maeda H, Nishida S, Hosen N, Tsuboi A, Oka Y and Sugiyama H. Translation elongation factor eEF2 is a novel tumor-associated antigen overexpressed in various types of cancers. Int Natl J Oncol 2014; 44: 1461-1469.	森 雅秀
Hirano H, Kimura A, Yamada H, Maeda H, Takeuchi Y, Susaki Y, Kobayashi R, Hayashi A, Ose N, <u>Yamaguchi T, Yokota S, Mori M</u> . Clinicopathological Characteristics of Pathological-Stage IA Lung Adenocarcinoma with a Solid-sheet Pattern Component. Acta Hyogo Med. 2014; 38(2): 69-74.	山口俊彦 横田総一郎 森 雅秀
Hirano H, Maeda H, <u>Yamaguchi T, Yokota S, Mori M</u> and Okimura A. Short Communication-Factors Related to Aggressiveness of Some Lung Carcinomas with Peculiar Histological Characteristics. JSM Clin Oncol Res 2(3):1022-1024.	山口俊彦 横田総一郎 森 雅秀

<p>Hirano H, Maeda H, Takeuchi Y, Susaki Y, Kobayashi R, Hayashi A, Ose N, <u>Yamaguchi T, Yokota S, Mori M</u>. Lymphatic invasion of micropapillary cancer cells is associated with a poor prognosis of pathological stage IA lung adenocarcinomas. Micropapillary adenocarcinoma, Oncol Lett. 2014 Sep;8(3):1107–1111. Epub 2014 Jun 25.</p>	<p>山口俊彦 横田総一郎 森 雅秀</p>
<p>Hirano H, <u>Niinaka M</u>, Kuroyama M, <u>Satomi A, Yokota S, Yamaguchi T, Mori M</u>. Pleomorphic and Desmoplastic Malignant Mesotheliomas and a Malignant Mesothelioma with Osseous and Cartilaginous Differentiation: Case Reports. J Cancer Biol Res 2014 2(1): 1024–1032.</p>	<p>里見明俊 新中 学 山口俊彦 横田総一郎 森 雅秀</p>
<p>Hirano H, Maeda H, Takeuchi Y, Susaki Y, Kobayashi R, Hayashi A, Ose N, <u>Ninaka M, Yamaguchi T, Yokota S, Mori M</u>. Immunohistochemical analysis of p–stage I large cell neuroendocrine carcinoma of the lung: Analysis of adhesion molecules and proliferative activity. J Cancer Biol Res 2014;2(1);1034–1040.</p>	<p>新中 学 山口俊彦 横田総一郎 森 雅秀</p>
<p>Hattori Y, Satouchi M, Shimada T, Urata Y , <u>Yoneda T, Mori M</u>, Nishimura T, Sunadome H, Kumagai T, Imamura, Fujita S, Kaji R, Hata A, Tachihara M, Morita S, Negoro S. A phase 2 study of bevacizumab in combination with carboplatin andpaclitaxel in patients with non-squamous non-small-cell lung cancerharboring mutations of epidermal growth factor receptor (EGFR) afterfailing first-line EGFR-tyrosine kinase inhibitors (HANSHIN Oncology Group 0109). Lung Cancer. 2015 Feb;87(2):136–140. doi: 10.1016/j.lungcan.2014.12.007. Epub 2014 Dec 19.</p>	<p>米田 勉 森 雅秀</p>